



Decay induced de-chelation of positron-emitting electron-capture daughters and its use in preclinical PET

Severin, Gregory; Kristensen, L.K. ; Fonslet, Jesper; Jensen, Andreas Tue Ingemann; Nielsen, C.H. ; Elema, Dennis Ringkjøbing; Johnston, K. ; Köster, U.

Published in:
Book of Abstracts. ISOLDE Workshop and Users meeting 2015

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Severin, G., Kristensen, L. K., Fonslet, J., Jensen, A. T. I., Nielsen, C. H., Elema, D. R., Johnston, K., & Köster, U. (2015). Decay induced de-chelation of positron-emitting electron-capture daughters and its use in preclinical PET. In *Book of Abstracts. ISOLDE Workshop and Users meeting 2015*

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Decay induced de-chelation of positron-emitting electron-capture daughters and its use in preclinical PET.

G.W. Severin¹, L.K. Kristensen², J. Fonslet¹, A.I. Jensen¹, C.H. Nielsen², D.R. Elema¹, K. Johnston³, and U. Köster⁴

¹*Hevesy Laboratory, Center for Nuclear Technologies at the Technical University of Denmark, Frederiksborgvej 399, 4000 Roskilde, Denmark.*

²*Department of Clinical Physiology, Nuclear Medicine & PET and Cluster for Molecular Imaging, Rigshospitalet and University of Copenhagen, Blegdamsvej 3, 2200 Copenhagen, Denmark.*

³*ISOLDE, CERN, CH-1211 Geneva 23, Switzerland*

⁴*Institut Laue-Langevin, 71 avenue des Martyrs, F-38042 Grenoble Cedex 9, France*

Abstract

The purpose of this study was to assess the value of $^{140}\text{Nd}/^{140}\text{Pr}$ and $^{134}\text{Ce}/^{134}\text{La}$ as *in vivo* PET generators through kinetic evaluation of the pairs with somatostatin analogues in the neuroendocrine tumor-based somatostatin receptor 2 (sst₂) system. By employing a known sst₂ internalizing vector, DOTATATE, and a known non-internalizing vector, DOTALM3, we attempted to determine *in vivo* diffusion kinetics of a freed positron-emitting daughter $^{140}\text{Pr}^{3+}$ in tumor-bearing mice.

Methods: ^{140}Nd and ^{134}Ce were produced by proton-induced spallation of a tantalum target at ISOLDE, electromagnetically separated, and implanted in a thin zinc layer on gold foils. After dissolving the zinc, the products were chemically separated, reacted with sst₂ targeting vectors DOTATATE or DOTALM3, and the ^{140}Nd labeled vectors were i.v. injected into dual-flank H727 xenograft bearing mice (n = 8 for each tracer). PET scans were taken at 1, 3, and 16 hours post injection. Following the last image, the animals were euthanized, and then imaged intact at 30 min post-mortem. The differences between the 16 h PET scan and the post-mortem scan were used to study the diffusion behavior of $^{140}\text{Pr}^{3+}$ following the parent ^{140}Nd EC decay. Three additional mice were scanned under the same protocol after injection of $^{140}\text{Nd}^{3+}$ in HEPES-buffered isotonic saline to investigate the free ion distributions in the tumor model.

Results: In total 950 MBq of ^{140}Nd and 140 MBq of ^{134}Ce were collected at ISOLDE and shipped to Hevesy Lab. ^{140}Nd reactions with DOTATATE and DOTALM3 were efficient at 5 MBq/nmol (n = 2 each). ^{134}Ce labeling was inefficient, and could only be achieved with receptor-saturating levels of the vectors, thereby precluding their use *in vivo*. The *in vivo* scans showed only a small difference in the tumor PET signal between pre- and post-mortem scans, with a slight increase in tumor signal post-mortem when DOTALM3 was used. Non-targeted organs, however, showed interesting source and sink behaviors illuminating some properties of the renal and hepatic interaction with neodymium and praseodymium.

Conclusion: Based upon the results of this study we conclude that ^{140}Nd imaging in preclinical models might be possible without designing an electron-capture-dislocation resistant chelate. In such cases, the imaging protocol established here is a useful test to determine how ^{140}Nd PET is altered by diffusion. Further, with careful experimental design it may be possible to exploit the diffusion effects to observe biological phenomena such as vector internalization.